

# Wabeek Lake 2021 Water Quality Report

A publication of the Wabeek Lake Improvement Board

**Wabeek Lake Improvement Board**  
4200 Telegraph Road  
Bloomfield Hills, MI 48303-0489

Dani Walsh, Chair  
Bloomfield Township Supervisor

Martin Brook, Secretary & Treasurer  
Bloomfield Township Clerk

Jay Shah  
Wabeek Lake Riparian

Chuck Moss  
Oakland County Commissioner

Jacy Garrison  
Oakland County Water Resources  
Commissioner's Office

This report provides background information on lake water quality and a discussion of Wabeek Lake 2021 sampling results.

Lakes can be classified into three broad categories based on their productivity or ability to support plant and animal life. The three basic lake classifications are oligotrophic, mesotrophic, and eutrophic.


*Oligotrophic* lakes are generally deep and clear with little aquatic plant growth. These lakes maintain sufficient dissolved oxygen in the cool, deep bottom waters during late summer to support cold water fish such as trout and whitefish.

*Eutrophic* lakes have poor clarity, and support abundant aquatic plant growth. In deep eutrophic lakes, the cool bottom waters usually contain little or no dissolved oxygen. Therefore, these lakes can only support warm water fish such as bass and pike.

Lakes that fall between the two extremes of oligotrophic and eutrophic are called *mesotrophic* lakes.

Under natural conditions, most lakes will ultimately evolve to a eutrophic state as they gradually fill with sediment and organic matter transported to the lake from the surrounding watershed.

As the lake becomes shallower, the process accelerates. When aquatic plants become abundant, the lake slowly begins to fill in as sediment and decaying plant matter accumulate on the lake bottom. Eventually, terrestrial plants become established and the lake is transformed to a marshland. The natural lake aging process can be greatly accelerated if excessive amounts of sediment and nutrients (which stimulate aquatic plant growth) enter the lake from the surrounding watershed. Because these added inputs are usually associated with human activity, this accelerated lake aging process is often referred to as *cultural eutrophication*.



Lake trophic states.

Environmental Consultant  
Progressive AE

Herbicide Applicator  
Aqua-Weed Control

# Trophic State Indicators

For more information, visit:  
[michiganlakeinfo.com/trophic-state](http://michiganlakeinfo.com/trophic-state)

2

Key parameters used to evaluate a lake's productivity or trophic state include total phosphorus, chlorophyll-*a*, and Secchi transparency.

*Phosphorus* is the nutrient that most often stimulates excessive growth of aquatic plants and causes premature lake aging. By measuring phosphorus levels, it is possible to gauge the overall health of a lake.


*Chlorophyll-a* is a pigment that imparts the green color to plants and algae. A rough estimate of the quantity of algae present in the water column can be made by measuring the amount of chlorophyll-*a* in the water column.

A *Secchi disk* is a round, black and white, 8-inch disk that is used to estimate water clarity. Generally, it has been found that plants can grow to a depth of about twice the Secchi disk transparency.

Generally, as phosphorus inputs to a lake increase, algae growth and chlorophyll-*a* increase and Secchi transparency decreases.

pH is a measure of the amount of acid or base in the water. The pH scale ranges from 0 (acidic) to 14 (alkaline or basic) with neutrality at 7. The pH of most lakes in the Upper Midwest ranges from 6.5 to 9.0.


The alkalinity of most lakes in the Upper Midwest is within the range of 23 to 148 milligrams per liter, or parts per million, as calcium carbonate.



A Secchi disk measures water clarity.

## TROPHIC CLASSIFICATION CRITERIA

Lake Classification	Total Phosphorus (µg/L) <sup>1</sup>	Chlorophyll- <i>a</i> (µg/L) <sup>1</sup>	Secchi Transparency (feet)
Oligotrophic	Less than 10	Less than 2.2	Greater than 15.0
Mesotrophic	10 to 20	2.2 to 6.0	7.5 to 15.0
Eutrophic	Greater than 20	Greater than 6.0	Less than 7.5



In 2021, samples were collected on July 28 at the surface, mid-depth and bottom from the deep basin of Wabek Lake.

<sup>1</sup> µg/L = micrograms per liter = parts per billion.

## Temperature and Dissolved Oxygen

Temperature and dissolved oxygen strongly influence lake water quality and are very important to a lake's fishery.

For more information, visit:  
[michiganlakeinfo.com/turnover-and-stratification](http://michiganlakeinfo.com/turnover-and-stratification)


### Temperature

*Temperature* is important in determining the type of organisms that may live in a lake. For example, trout prefer temperatures below 68°F. Temperature also determines how water mixes in a lake. As the ice cover breaks up on a lake in the spring, the water temperature becomes uniform from the surface to the bottom. This period is referred to as *spring turnover* because water mixes throughout the entire water column. As the surface waters warm, they are underlain by a colder, more dense layer of water. This process is called thermal stratification. In deeper lakes during summer there are three distinct layers. This is referred to as *summer stratification*. Once thermal stratification occurs, there is little mixing of the warm surface waters with the cooler bottom waters. The transition layer that separates these layers is referred to as the *thermocline*. The thermocline is characterized as the zone where temperature drops rapidly with depth. As fall approaches, the warm surface waters begin to cool and become more dense. Eventually, the surface temperature drops to a point that allows the lake to undergo complete mixing. This period is referred to as *fall turnover*. As the season progresses and ice begins to form on the lake, the lake may stratify again. However, during *winter stratification*, the surface waters (at or near 32°F) are underlain by slightly warmer water (about 39°F). This is sometimes referred to as *inverse stratification* and occurs because water is most dense at a temperature of about 39°F. As the lake ice melts in the spring, these stratification cycles are repeated. These stratification cycles occur in deep lakes but not in shallow lakes or ponds. Lakes that are about 15 to 30 feet deep may stratify and destratify with storm events several times during the year. Wabek Lake is deep enough to stratify during the summer months and likely stratifies as well under ice cover during the winter.

### Dissolved Oxygen

An important factor influencing lake water quality is the quantity of *dissolved oxygen* in the water column. The major inputs of dissolved oxygen to lakes are the atmosphere and photosynthetic activity by aquatic plants. An oxygen level of about 5 mg/L (milligrams per liter, or parts per million) is required to support warm-water fish. In lakes deep enough to exhibit thermal stratification, oxygen levels are often reduced or depleted below the thermocline once the lake has stratified. This is because deep water is cut off from plant photosynthesis and the atmosphere, and oxygen is consumed by bacteria that use oxygen as they decompose organic matter (plant and animal remains) at the bottom of the lake. Bottom-water oxygen depletion is a common occurrence in eutrophic and some mesotrophic lakes. Thus, eutrophic and most mesotrophic lakes cannot support cold-water fish because the cool, deep water (that the fish require to live) does not contain sufficient oxygen.

3



**Seasonal thermal stratification.** Stratification cycles occur in deep lakes but not in shallow lakes or ponds.

# Wabeek Lake Water Quality

4

Water quality data collected from the deep basin of Wabeek Lake on July 28, 2021 are summarized in the tables below. At the time of sampling, Wabeek Lake was thermally stratified and devoid of oxygen in the deep portion of the lake. The elevated phosphorus levels at the lake bottom are the result of phosphorus release from the deep water sediments in the lake which commonly occurs in the absence of oxygen. pH and alkalinity are within a normal range for Michigan lakes. Data collected in 2021 are generally consistent with historical water quality data for Wabeek Lake. The surface water phosphorus concentration, Secchi transparency, and chlorophyll-a data indicate Wabeek Lake is mesotrophic. Data also indicates that Wabeek Lake will support warm water fish species, such as bluegill and largemouth bass.

**TABLE 1**  
**WABEEK LAKE**  
**JULY 28, 2021 DEEP BASIN WATER QUALITY DATA**

Date	Station	Sample Depth (feet)	Temperature (°F)	Dissolved Oxygen (mg/L) <sup>1</sup>	Total Phosphorus (µg/L) <sup>2</sup>	pH (S.U.) <sup>3</sup>	Total Alkalinity (mg/L CaCO <sub>3</sub> ) <sup>4</sup>
28-Jul-21	1	1	80	8.7	19	8.7	118
28-Jul-21	1	13	73	2.3	40	8.3	130
28-Jul-21	1	25	48	0.0	127	7.6	191

**TABLE 2**  
**WABEEK LAKE**  
**JULY 28, 2021 SURFACE WATER QUALITY DATA**

Date	Station	Secchi Transparency (feet)	Chlorophyll-a (µg/L) <sup>2</sup>
28-Jul-21	1	8.5	4

1 mg/L = milligrams per liter = parts per million.

2 µg/L = micrograms per liter = parts per billion.

3 S.U. = standard units.

4 mg/L CaCO<sub>3</sub> = milligrams per liter as calcium carbonate.